arXiv:1301.5729v2 [math.GT] 14 Mar 2013

LEFT-ORDERABLE, NON–L–SPACE SURGERIES ON KNOTS

KIMIHIKO MOTEGI AND MASAKAZU TERAGAITO

Abstract. A Dehn surgery is said to be *left-orderable* if the resulting manifold of the surgery has the left-orderable fundamental group, and a Dehn surgery is called an *L–space surgery* if the resulting manifold of the surgery is an L–space. We will focus on left-orderable, non–L–space surgeries on knots in the 3–sphere. Once we have a knot with left-orderable surgeries, the "periodic construction" enables us to provide infinitely many knots with left-orderable, non–L–space surgeries. We apply the construction to present infinitely many hyperbolic knots on each of which every nontrivial surgery is a left-orderable, non–L–space surgery.

1. Introduction

A nontrivial group G is said to be *left-orderable* if there exists a strict total ordering $<$ on its elements such that $g < h$ implies $fg < fh$ for all elements $f, g, h \in G$. The left-orderability of fundamental groups of 3–manifolds has been studied by Boyer, Rolfsen and Wiest [4]. In particular, they prove that the fundamental group of a P^2–irreducible 3–manifold is left-orderable if and only if it has an epimorphism to a left-orderable group [4, Theorem 1.1(1)]. Since the infinite cyclic group $\mathbb{Z}$ is left-orderable, a P^2–irreducible 3–manifold with first Betti number $b_1 \geq 1$ has a left-orderable fundamental group. One obstruction for G being left-orderable is an existence of torsion elements in G. Thus, for instance, lens spaces cannot have left-orderable fundamental groups. It is interesting to characterize rational homology 3–spheres whose fundamental groups are left-orderable. Examples suggest that there exists a correspondence between 3–manifolds whose fundamental groups are left-orderable and L–spaces which appear in the Heegaard Floer homology theory [36, 37]. Recall that a rational homology 3–sphere Y is called an *L–space* if the rank of its Heegaard Floer homology $\widehat{HF}(Y)$ coincides with $|H_1(Y;\mathbb{Z})|$. The name stems from the fact that lens spaces are L–spaces.

The following conjecture is formulated by Boyer, Gordon and Watson [3].

The first author has been partially supported by JSPS Grants–in–Aid for Scientific Research (C) (No.21540098), The Ministry of Education, Culture, Sports, Science and Technology, Japan and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2012.
The second author has been partially supported by JSPS Grant-in-Aid for Scientific Research (C), 22540088, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

2010 *Mathematics Subject Classification.* Primary 57M25, 57N10, 20F60 Secondary 06F15

Conjecture 1.1. *An irreducible rational homology 3–sphere is an L–space if and only if its fundamental group is not left-orderable.*

In [3] the conjecture is verified for geometric, non-hyperbolic 3–manifolds and the 2–fold branched covers of non-splitting alternating links. See also [13, 19, 40] for related results.

A useful way to construct rational homology 3–spheres is Dehn surgery on knots in the 3–sphere S^3. For any knot K in S^3 the exterior $E(K) = S^3 - \mathrm{int}N(K)$ has the left-orderable fundamental group, and the longitudinal surgery (i.e. 0–surgery) on K yields a 3–manifold with left-orderable fundamental group [4, Theorem 1.1]. On the other hand, the result $K(r)$ of r–Dehn surgery may not have such a fundamental group if $r \neq 0$; see [10]. A Dehn surgery is said to be *left-orderable* if the resulting manifold of the surgery has the left-orderable fundamental group, and a Dehn surgery is called an *L–space surgery* if the resulting manifold of the surgery is an L–space. For the figure-eight knot K it is known that every nontrivial surgery is a non–L–space surgery [38, 39], and thus it is expected that every nontrivial surgery on K is left-orderable. Boyer, Gordon and Watson [3] show that $\pi_1(K(r))$ is left-orderable if $-4 < r < 4$, and Clay, Lidman and Watson [7] verify that it is left-orderable for $r = \pm 4$. For further related results, see [8, 14, 15, 19, 45, 47, 48] which give supporting evidence for Conjecture 1.1.

In the present note, we will focus on left-orderable, non–L–space surgeries on knots in S^3. First we show an "inheritance" property of left-orderability: The fundamental groups of 3–manifolds obtained by Dehn surgeries on a periodic knot K inherit the left-orderability from those of 3–manifolds obtained by Dehn surgeries on the factor knot $\overline{K}$; see Theorem 2.1. Then based on the inheritance property, we introduce a "periodic construction" (Corollary 4.1) which enables us to provide infinitely many hyperbolic knots having left-orderable, non–L–space surgeries from a given knot with left-orderable surgeries. See Corollary 4.1 for the precise statement. In Sections 5, 6 we will give some examples illustrating how the periodic construction works. In Section 7 we will apply the "periodic construction", together with the "satellite construction" ([9, Proposition 4.1]) to prove the following results.

Theorem 1.2. *There exist infinitely many hyperbolic knots K each of which enjoys the following properties.*

(1) *$K(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$.*
(2) *$K(r)$ is not an L–space for all $r \in \mathbb{Q}$.*
(3) *$\pi_1(K(r))$ is left-orderable for all $r \in \mathbb{Q}$.*

Theorem 1.3. *For a given integer $N > 0$, there exists a knot K which enjoys the following properties.*

(1) *The genus and the tunnel number of K are both greater than N.*
(2) *The Heegaard genus of $K(r)$ is greater than N for all but finitely many r.*
(3) *$K(r)$ is not an L–space for all $r \in \mathbb{Q}$.*
(4) *$\pi_1(K(r))$ is left-orderable for all $r \in \mathbb{Q}$.*

2. Left-orderable surgeries on periodic knots

A knot K in S^3 is called a *periodic knot* with period p if there is an orientation preserving diffeomorphism $f : S^3 \to S^3$ such that $f(K) = K$, $f^p = id$ $(p > 1)$, $\mathrm{Fix}(f) \neq \emptyset$, and $\mathrm{Fix}(f) \cap K = \emptyset$, where $\mathrm{Fix}(f)$ is the set of fixed points of f. By the positive answer to the Smith conjecture [31], f is a rotation of S^3 about the unknotted circle $C = \mathrm{Fix}(f)$. So by taking the quotient $S^3/\langle f \rangle$, we obtain the *factor knot* $\overline{K} = K/\langle f \rangle$ and the unknotted circle $\overline{C} = C/\langle f \rangle$ in $S^3 = S^3/\langle f \rangle$. We often call C the *axis* and $\overline{C}$ the *branch circle*. Since K is connected, the linking number $lk(\overline{C}, \overline{K})$ and the period p are relatively prime. Note that if the periodic knot K is unknotted, then the equivariant loop theorem [27] implies that $K \cup C$ is the Hopf link and $\overline{K} \cup \overline{C}$ is also the Hopf link. To exclude such a trivial case, in the following we consider nontrivial periodic knots.



Figure 2.1. A periodic knot and its factor knot

The next theorem asserts that the fundamental groups of 3–manifolds obtained by Dehn surgeries on the periodic knot K inherit the left-orderability from those of 3–manifolds obtained by Dehn surgeries on the factor knot $\overline{K}$. For a subset $\mathcal{S} \subset \mathbb{Q}$ and a positive integer p, we denote by $p\mathcal{S}$ the subset $\{pr \mid r \in \mathcal{S}\} \subset \mathbb{Q}$. Note that if $\mathcal{S} = \mathbb{Q}$, then $p\mathcal{S} = \mathbb{Q}$.

Theorem 2.1. *Let K be a nontrivial knot in S^3 with cyclic period p, and let $\overline{K}$ be its factor knot. Suppose that $\pi_1(\overline{K}(r))$ is left-orderable for $r \in \mathcal{S}$. Then $\pi_1(K(s))$ is left-orderable for $s \in p\mathcal{S}$.*

Proof of Theorem 2.1. Let $f : S^3 \to S^3$ be an orientation preserving diffeomorphism giving the cyclic period p of K with the axis $C = \mathrm{Fix}(f)$ and the factor knot $\overline{K} = K/\langle f \rangle$. Take an $\langle f \rangle$–invariant tubular neighborhood $N(K)$ of K. Let $N(\overline{K})$

be the quotient $N(K)/\langle f\rangle$. In the following $E(K) = S^3 - \mathrm{int}N(K)$ and $E(\overline{K}) = E(K)/\langle f\rangle = S^3 - \mathrm{int}N(\overline{K})$. Denote by (μ, λ) (resp. $(\overline{\mu}, \overline{\lambda})$) a preferred meridian-longitude pair of $\pi_1(\partial N(K))$ (resp. $\pi_1(\partial N(\overline{K}))$). We can choose a simple closed curve representing the preferred longitude λ which is invariant under $\langle f\rangle$; see [11].

Let $\pi : E(K) \to E(\overline{K})$ be the cyclic branched covering branched along $\overline{C} = C/\langle f\rangle$.

Lemma 2.2. *The branched cover* $\pi : E(K) \to E(\overline{K})$ *is extended to a branched cover* $\pi' : K(\frac{m}{n}) \to \overline{K}(\frac{m}{pn})$.

Proof of Lemma 2.2. Let $\pi_* : \pi_1(E(K)) \to \pi_1(E(\overline{K}))$ be the homomorphism induced by π. Then $\pi_*|_{\pi_1(\partial E(K))} : \pi_1(\partial E(K)) \to \pi_1(\partial E(\overline{K}))$ sends μ to $\overline{\mu}$, and λ to $p\overline{\lambda}$. Hence $\pi_*|_{\pi_1(\partial E(K))}(m\mu+n\lambda) = m\overline{\mu}+pn\overline{\lambda} = (m,\ p)\{\frac{m}{(m,\ p)}\overline{\mu}+\frac{pn}{(m,\ p)}\overline{\lambda}\}$. Then we can extend $\pi : E(K) \to E(\overline{K})$ to $\pi' : K(\frac{m}{n}) = E(K)\cup V \to \overline{K}(\frac{m}{pn}) = E(\overline{K})\cup\overline{V}$, where V, $\overline{V}$ are filled solid tori. If $(m,p) \geq 2$, then π' branches along the core of the filled solid torus $\overline{V}$ as well as $\overline{C}$. □(Lemma 2.2)

Thus we have a commutative diagram:

$$\begin{tikzcd}
E(K) \arrow[r, "\pi"] \arrow[d, "\text{Dehn filling}"'] & E(\overline{K}) \arrow[d, "\text{Dehn filling}"] \\
K(\frac{m}{n}) \arrow[r, "\pi'"'] & \overline{K}(\frac{m}{pn})
\end{tikzcd}$$

Assume that $\frac{m}{pn} \in \mathcal{S}$, i.e. $\overline{K}(\frac{m}{pn})$ has the left-orderable fundamental group. Let us prove that $K(\frac{m}{n})$ has also the left-orderable fundamental group.

Lemma 2.3. $K(\frac{m}{n})$ *is irreducible.*

Proof of Lemma 2.3. Suppose for a contradiction that $K(\frac{m}{n})$ is reducible. Since K is a nontrivial periodic knot, K is cabled and $\frac{m}{n}$ is the cabling slope [25, 16, 17]. First we assume that K is a torus knot. Then $E(K)$ has a unique Seifert fibration (up to isotopy). Following [26, Theorem 2.2], we choose a Seifert fibration of $E(K)$ which is preserved by f. If C is not a fiber, we take a regular fiber t intersecting C. Since f fixes a point in $t\cap C$, $f(t) = t$ and f reverses the orientation of t. This then implies that f reverses the orientation of K, and hence C intersects K in exactly two points, a contradiction. Thus C is a fiber in the $\langle f\rangle$–invariant Seifert fibration of $E(K)$. Since a regular fiber is knotted in S^3, C is one of two exceptional fibers in $E(K)$. Then the quotient $E(\overline{K}) = E(K)/\langle f\rangle$ has also a Seifert fibration induced from that of $E(K)$ and thus $\overline{K}$ is a torus knot; the surgery slope $\frac{m}{pn}$ on $\partial E(\overline{K})$ is the fiber slope. Since $\frac{m}{pn}$ is the fiber (i.e. cabling) slope, $\pi_1(\overline{K}(\frac{m}{pn}))$ has a nontrivial torsion, contradicting the left-orderability of $\pi_1(\overline{K}(\frac{m}{pn}))$.

Next assume that K is an (x, y)–cable in a knotted solid torus W, where $y \geq 2$. By the $\langle f \rangle$–invariant version ([26, Theorem 8.6]) of the torus decomposition theorem [20, 21], we may assume that f leaves a companion solid torus W invariant. First we note that $W \cap C = \emptyset$. For otherwise, $f|_{\partial W}$ has fixed points and hence it is an involution, and f reverses the orientation of an $\langle f \rangle$–invariant core of W. Hence it also reverses the orientation of K (which has the winding number $y \geq 2$ in W). This then implies that C intersects K in exactly two points, a contradiction. Thus $W \subset S^3 - C$. We denote the quotient $W/\langle f \rangle$ by $\overline{W}$. We may assume that the cable space $W - \mathrm{int}N(K)$ has a Seifert fibration preserved by f. Then $\overline{W} - \mathrm{int}N(\overline{K}) = (W - \mathrm{int}N(K))/\langle f \rangle$ has an induced Seifert fibration in which a regular fiber on $\partial N(\overline{K})$ represents the surgery slope $\frac{m}{pn}$. This implies that the result of $\frac{m}{pn}$–surgery of $\overline{W}$ along $\overline{K}$, and hence $\overline{K}(\frac{m}{pn})$, has a nontrivial lens space summand whose fundamental group has order $y \geq 2$. Since $\pi_1(\overline{K}(\frac{m}{pn}))$ has a nontrivial torsion, it cannot be left-orderable, contradicting the assumption. □(Lemma 2.3)

The above diagram induces the commutative diagram of fundamental groups below.

$$\begin{tikzcd}
\pi_1(E(K)) \arrow[r, "\pi_*"] \arrow[d] & \pi_1(E(\overline{K})) \arrow[d] \\
\pi_1(K(\frac{m}{n})) \arrow[r, "\pi'_*"'] & \pi_1(\overline{K}(\frac{m}{pn}))
\end{tikzcd}$$

Lemma 2.4. $\pi_* : \pi_1(K(\frac{m}{n})) \to \pi_1(\overline{K}(\frac{m}{pn}))$ *is surjective.*

Proof of Lemma 2.4. Choose a point $x \in C = \mathrm{Fix}(f)$ (resp. $\pi(x) \in \overline{C}$) as a base point of $\pi_1(E(K))$ (resp. $\pi_1(E(\overline{K}))$. We take obvious meridians $\overline{\mu}_i$ of $\overline{K}$ which are generators of $\pi_1(E(\overline{K}), \pi(x))$ (with respect to the Wirtinger presentation of $\pi_1(E(\overline{K}), \pi(x))$). Then their lifts $\mu_i \in \pi_1(E(K))$ satisfy $\pi_*(\mu_i) = \overline{\mu}_i$, and hence $\pi_* : \pi_1(E(K)) \to \pi_1(E(\overline{K}))$ is an epimorphism. Since vertical homomorphisms are also epimorphisms, $\pi'_* : \pi_1(K(\frac{m}{n})) \to \pi_1(\overline{K}(\frac{m}{pn}))$ is also an epimorphism. □(Lemma 2.4)

By Lemma 2.3 $K(\frac{m}{n})$ is irreducible, and by Lemma 2.4 we have an epimorphism from $\pi_1(K(\frac{m}{n}))$ to the left-orderable group $\pi_1(\overline{K}(\frac{m}{pn}))$. Then it follows from [4, Theorem 1.1(1)] that $\pi_1(K(\frac{m}{n}))$ is also left-orderable. Thus if $\pi_1(\overline{K}(r))$ is left-orderable for $r = \frac{m}{pn} \in \mathcal{S}$, then $\pi_1(K(s))$ is also left-orderable for $s = pr = \frac{m}{n} \in p\mathcal{S}$. □(Theorem 2.1)

For a given knot K, denote by pK the connected sum of p–copies of K. Then obviously pK is a periodic knot with the factor knot K. Thus Theorem 2.1 immediately implies:

Corollary 2.5. *Let K be a knot in S^3 such that $\pi_1(K(r))$ is left-orderable for $r \in \mathcal{S}$. Then $\pi_1(pK(s))$ is left-orderable for any $s \in p\mathcal{S}$.*

3. L–SPACE SURGERIES ON PERIODIC KNOTS

In [34, 35] Ni proves that if a knot K in S^3 has an L–space surgery, then K is a fibered knot, i.e. $E(K)$ has a fibering over the circle. For a periodic knot K, the next theorem gives a necessary condition on the factor knot for K having an L–space surgery.

Theorem 3.1. *Let K be a periodic knot in S^3 with the axis C, and let $\overline{K}$ be its factor knot with the branch circle $\overline{C}$. Suppose that K has an L–space surgery. Then $E(\overline{K})$ has a fibering over the circle with a fiber surface $\overline{S}$ such that $|\overline{C} \cap \overline{S}|$ equals the algebraic intersection number between $\overline{C}$ and $\overline{S}$, i.e. the linking number $lk(\overline{C}, \overline{K})$.*

In particular, we have:

Corollary 3.2. *Let K be a periodic knot with the factor knot $\overline{K}$. If $\overline{K}$ is not fibered, then $K(r)$ is not an L–space for all $r \in \mathbb{Q}$.*

Proof of Theorem 3.1. Let $f : S^3 \to S^3$ be an orientation preserving diffeomorphism of finite order satisfying $f(K) = K$. Note that $C = \mathrm{Fix}(f)$, $\overline{K} = K/\langle f \rangle$ and $\overline{C} = C/\langle f \rangle$. Let $N(K)$ be an $\langle f \rangle$–invariant tubular neighborhood of K.

Assume that K has an L–space surgery. Then Ni [34, Corollary 1.3] ([35]) proves that $E(K) = S^3 - \mathrm{int}N(K)$ has a fibering over the circle. Following Proposition 6.1 in [12], we can isotope the fibering to a fibering preserved by the action of $\langle f \rangle$ so that the axis C is transverse to the fibers. Thus $E(\overline{K})$ inherits a fibering over the circle such that all the fibers transverse to the branch circle $\overline{C} = C/\langle f \rangle$. Let $\overline{S}$ be a fiber surface of $E(\overline{K})$. Since $\overline{C}$ intersects each fiber surface of the fibering of $E(\overline{K})$ transversely, $|\overline{C} \cap \overline{S}|$ coincides with the algebraic intersection number between $\overline{C}$ and $\overline{S}$, i.e. the linking number $lk(\overline{C}, \partial\overline{S})$, which equals the linking number $lk(\overline{C}, \overline{K})$. □(Theorem 3.1)

As Ni [34, 35] proves, the fiberedness of K is necessary for K having an L–space surgery. On the other hand, the periodicity of K itself also puts strong restrictions on 3–manifolds obtained by Dehn surgeries on K. For instance, if a periodic knot with period $p > 2$ has a finite surgery, then K is a torus knot or a cable of a torus knot [29, Proposition 5.6]. So we would like to ask:

Question 3.3. Let K be a knot in S^3 with cyclic period $p > 2$ other than a torus knot, a cable of a torus knot. Then does K admit an L–space surgery?

4. Periodic constructions

Once we have a knot K such that $\pi_1(K(r))$ is left-orderable for $r \in \mathcal{S} \subset \mathbb{Q}$, the combination of Theorems 2.1 and 3.1 gives us the following effective method for providing yet further infinitely many knots K' in S^3 for each of which $K'(s)$ is not an L–space and has the left-orderable fundamental group whenever s belongs to a subset $\mathcal{S}' \subset \mathbb{Q}$.

Corollary 4.1 (periodic construction). *Let $\overline{K}$ be a knot in S^3 such that $\pi_1(\overline{K}(r))$ is left-orderable for $r \in \mathcal{S}$. Take an unknotted circle $\overline{C}$ disjoint from $\overline{K}$ with linking number $lk(\overline{C}, \overline{K}) = l$. If $\overline{K}$ is a fibered knot, then we choose $\overline{C}$ so that any fiber surface (i.e. minimal genus Seifert surface) $\overline{S}$ satisfies the condition $|\overline{C} \cap \overline{S}| > lk(\overline{C}, \overline{K})$. Let p be an integer such that $p \geq 2$ and $(p, l) = 1$. Take the p–fold cyclic branched cover of S^3 branched along $\overline{C}$ to obtain a periodic knot $K_{\frac{p}{C}}$ which is the preimage of $\overline{K}$. Then $K_{\frac{p}{C}}$ enjoys the following properties:*

1. *$K_{\frac{p}{C}}(s)$ is not an L–space for all $s \in \mathbb{Q}$.*
2. *$\pi_1(K_{\frac{p}{C}}(s))$ is left-orderable whenever $s \in p\mathcal{S}$.*

If $\overline{K}$ is a trivial knot, then the left-orderability of $\pi_1(\overline{K}(r))$ implies that $r = 0$, i.e. $\mathcal{S} = \{0\}$, and hence $p\mathcal{S} = \{0\}$. So we will apply Corollary 4.1 in the case where $\overline{K}$ is nontrivial.

Remark 4.2. We denote the genus of a knot k in S^3 by $g(k)$. For $\overline{K}$ and $K_{\frac{p}{C}}$, we have $g(K_{\frac{p}{C}}) \geq pg(\overline{K})$ [33, Theorem 3.2].

When we apply Corollary 4.1 for a given nontrivial knot $\overline{K}$, there are infinitely many choices for $\overline{C}$, and we can expect that in most cases, $K_{\frac{p}{C}}$ are hyperbolic knots and $K_{\frac{p}{C}}(s)$ are hyperbolic 3–manifolds. In fact, we can prove the following.

Theorem 4.3. *For a given nontrivial knot $\overline{K}$ in S^3, we have the following.*

1. *There are infinitely many unknotted circles $\overline{C}$ such that $\overline{K} \cup \overline{C}$ is a hyperbolic link.*
2. *If $\overline{K} \cup \overline{C}$ is a hyperbolic link and $p > 2$, then $K_{\frac{p}{C}}$ is a hyperbolic knot, and $K_{\frac{p}{C}}(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$.*
3. *Assume that $p > 2$ and $\overline{C_i}$ $(i = 1, 2)$ is an unknotted circle such that $lk(\overline{C_i}, \overline{K})$ and p are relatively prime, and $\overline{K} \cup \overline{C_i}$ is a hyperbolic link. If $K_{\frac{p}{C_1}}$ and $K_{\frac{p}{C_2}}$ are isotopic in S^3, then $\overline{K} \cup \overline{C_1}$ and $\overline{K} \cup \overline{C_2}$ are isotopic.*

Proof of Theorem 4.3. (1) The following argument is due to Aitchison [1]. Arrange $\overline{K}$ as a closed n–braid for some integer n. Then introduce $(n-1)$–strands

C_i $(i = 1, \dots, n-1)$ between the n–strands of the original braid so that the crossings introduced, together with the original crossings, are alternatively positive and negative. See Figure 4.1.



FIGURE 4.1. Insertion of $(n-1)$–strands

Then we arrange C_i as in Figure 4.2(i) so that the closed braid is a 2–component link consisting of $\overline{K}$ and an unknotted circle $\overline{C} = C_1 \cup \cdots \cup C_{n-1}$ and $\overline{K} \cup \overline{C}$ is a non-split prime alternating link [28, Theorem 1]. To see that there are infinitely many such $\overline{C}$, we add m–twist as in Figure 4.2(ii). Since the linking number is changed depending on the number m, we have infinitely many $\overline{C}$ with the desired property. It follows from [28, Corollary 2] that $\overline{K} \cup \overline{C}$ is either a torus link or a hyperbolic link. Since $\overline{K}$ is nontrivial, but $\overline{C}$ is trivial, the former cannot occur, and thus $\overline{K} \cup \overline{C}$ is a hyperbolic link.

(2) Assume for a contradiction that $K^p_{\overline{C}}$ is not hyperbolic. Then it is either a torus knot or a satellite knot. Let $f : S^3 \to S^3$ be the deck transformation of the p–fold cyclic branched cover given in Corollary 4.1, which is an orientation preserving diffeomorphism giving the cyclic period p of $K^p_{\overline{C}}$. In the following, we take an $\langle f \rangle$–invariant tubular neighborhood $N(K^p_{\overline{C}})$ and denote $S^3 - \mathrm{int}N(K^p_{\overline{C}})$ by $E(K^p_{\overline{C}})$. The preimage of the branch circle $\overline{C}$ is an unknotted circle $C = \mathrm{Fix}(f)$, which is contained in the interior of $E(K^p_{\overline{C}})$. Note also that $K^p_{\overline{C}}$ is a nontrivial knot. For otherwise, the equivariant loop theorem [27] implies that $K^p_{\overline{C}} \cup C$ is the Hopf link and $\overline{K} \cup \overline{C}$ is also the Hopf link, contradicting the nontriviality of $\overline{K}$.

Claim 4.4. *$K^p_{\overline{C}}$ is not a torus knot.*

Proof of Claim 4.4. Assume for a contradiction that $K^p_{\overline{C}}$ is a torus knot. Then $E(K^p_{\overline{C}})$ has a unique Seifert fibration up to isotopy. We choose a Seifert fibration of $E(K^p_{\overline{C}})$ which is preserved by f [26, Theorem 2.2]. Then the argument in the proof of Lemma 2.3 shows that C is one of two exceptional fibers in $E(K^p_{\overline{C}})$. Then the quotient $E(\overline{K}) - \mathrm{int}N(\overline{C}) = (E(K^p_{\overline{C}}) - \mathrm{int}N(C))/\langle f \rangle$ has also a Seifert fibration. Thus $S^3 - \mathrm{int}N(\overline{K} \cup \overline{C}) = E(\overline{K}) - \mathrm{int}N(\overline{C})$ is a Seifert fiber space, contradicting its hyperbolicity. □(Claim 4.4)



FIGURE 4.2. Arrangement of $C_1, \dots, C_{n-1}$; $n = 4$

Claim 4.5. *$K^{p}_{\overline{C}}$ is not a satellite knot.*

Proof of Claim 4.5. Suppose for a contradiction that $K^{p}_{\overline{C}}$ is a satellite knot. Then we have an $\langle f \rangle$–invariant torus decomposition of $E(K^{p}_{\overline{C}})$ [26, Theorem 8.6]. Let Σ be the invariant family of essential tori in $E(K^{p}_{\overline{C}})$.

Case (i). There is an essential torus $T \in \Sigma$ such that $f(T) = T$. Then T bounds an $\langle f \rangle$–invariant companion solid torus W containing $K^{p}_{\overline{C}}$. Note that $K^{p}_{\overline{C}}$ is not a core of W. We see that $W \cap C = \emptyset$, for otherwise $f|_{\partial W}$ has a fixed point and it is an involution, i.e. $(f|_{\partial W})^2$ is the identity map. By the classical Smith theory [44] f itself is an involution, contradicting the assumption. Thus W lies in $S^3 - C$. We may assume that $W \subset S^3 - \mathrm{int}N(C)$ for a small tubular neighborhood $N(C)$ of C. Since the core of W is not a core of $S^3 - \mathrm{int}N(C)$, $S^3 - \mathrm{int}N(K^{p}_{\overline{C}} \cup C)$ contains the $\langle f \rangle$–invariant essential torus $T = \partial W$. This then implies that $S^3 - \mathrm{int}N(\overline{K} \cup \overline{C})$ contains an essential torus $\partial W / \langle f \rangle$. This contradicts the hyperbolicity of $S^3 - \mathrm{int}N(\overline{K} \cup \overline{C})$.

Case (ii). For each $T \in \Sigma$, $f(T) \neq T$ (hence $f(T) \cap T = \emptyset$). Let us pick an essential torus $T \in \mathcal{T}$. Note that T is essential in $S^3 - \mathrm{int}N(K^{p}_{\overline{C}} \cup C)$. Then the image $\overline{T} \subset E(\overline{K} \cup \overline{C})$ of T by the covering projection is also essential. This contradicts the hyperbolicity of $S^3 - \mathrm{int}N(\overline{K} \cup \overline{C})$. $\square$(Claim 4.5)

It follows that $K^{p}_{\overline{C}}$ is a hyperbolic knot in S^3.

Since $K^{p}_{\overline{C}}$ is a hyperbolic knot with period $p > 2$, it follows from [30, Corollary 1.4] that $K^{p}_{\overline{C}}(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$, or $p = 3$, $r = 0$ and $g(K^{p}_{\overline{C}}) = 1$. Since $g(K^{p}_{\overline{C}}) \geq pg(\overline{K}) \geq p > 2$, the latter cannot occur. Hence $K^{p}_{\overline{C}}(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$ as desired.

(3) In the following, for notational simplicity, we write $K_i = K^{p}_{\overline{C_i}}$.

The assumption, together with (2), implies that K_i $(i = 1, 2)$ is a hyperbolic knot. Recall that K_i has an orientation preserving diffeomorphism f_i such that $f_i(K_i) = K_i$, $f_i^p = id$ and $\mathrm{Fix}(f_i) = C_i$. Note that $\overline{K} = K_i/\langle f_i\rangle$ and $\overline{C_i} = C_i/\langle f_i\rangle$. Suppose that K_1 and K_2 are isotopic in S^3. Then we have an orientation preserving diffeomorphism φ of S^3 such that $\varphi(K_1) = K_2$. Note that the orientation preserving diffeomorphism $f_2' = \varphi^{-1} \circ f_2 \circ \varphi$ of S^3 satisfies $f_2'(K_1) = K_1$ and gives also a cyclic period p for K_1. Let us put $C_2' = \varphi^{-1}(C_2)$. Then we see that $\mathrm{Fix}(f_2') = C_2'$. Since $\varphi \circ f_2' = f_2 \circ \varphi$, φ induces an orientation preserving diffeomorphism $\overline{\varphi}$: $S^3 = S^3/\langle f_2'\rangle \to S^3 = S^3/\langle f_2\rangle$ sending $K_1/\langle f_2'\rangle$ to $\overline{K} = K_2/\langle f_2\rangle$ and $C_2'/\langle f_2'\rangle$ to $\overline{C_2} = C_2/\langle f_2\rangle$.

Now the hyperbolic knot K_1 has two orientation preserving periodic diffeomorphisms f_1 and f_2' of period $p > 2$. Then [2, 2.1 Theorem (a)] shows that the pairwise isotopy classes $[f_1]$ and $[f_2']$ have order p in the symmetry group $\mathrm{Sym}(S^3, K_1)$. Furthermore, since K_1 is hyperbolic, $\mathrm{Sym}(S^3, K_1)$ is isomorphic to a finite cyclic group or a dihedral group [22, Theorems 10.5.3 and 10.6.2(2)]. This implies that subgroups $\langle [f_1]\rangle$ and $\langle [f_2']\rangle$ of order $p(> 2)$ coincide. Thus we have $[f_1] = [f_2']^k = [(f_2')^k]$ for some integer k $(1 \leq k \leq p-1)$ in $\mathrm{Sym}(S^3, K_1)$. Note that $(f_2')^k$ has also period p. Then it follows from [2, 2.1 Theorem (c)] that f_1 and $(f_2')^k$ are conjugate by a diffeomorphism g in $\mathrm{Diff}(S^3, K_1)$ which is isotopic to the identity, i.e. $(f_2')^k = g \circ f_1 \circ g^{-1}$ and $g(K_1) = K_1$. Note that $(f_2')^k$ leaves K_1 invariant and $K_1/\langle (f_2')^k\rangle = K_1/\langle f_2'\rangle$, and that $\mathrm{Fix}((f_2')^k) = C_2'$ and $C_2'/\langle (f_2')^k\rangle = C_2'/\langle f_2'\rangle$. For any $x \in C_1 = \mathrm{Fix}(f_1)$, we have $(f_2')^k(g(x)) = g(f_1(x)) = g(x)$, thus $g(x) \in \mathrm{Fix}((f_2')^k) = C_2'$, and hence $g(C_1) \subset C_2'$. Conversely if $x' \in C_2' = \mathrm{Fix}((f_2')^k)$, then we see that $g^{-1}(x') \in C_1$ and $x' \in g(C_1)$, hence $C_2' \subset g(C_1)$. Thus we have $g(C_1) = C_2'$. Therefore we have an orientation preserving diffeomorphism $\overline{g} : S^3 = S^3/\langle f_1\rangle \to S^3 = S^3/\langle (f_2')^k\rangle = S^3/\langle f_2'\rangle$ sending $\overline{K} = K_1/\langle f_1\rangle$ to $K_1/\langle (f_2')^k\rangle = K_1/\langle f_2'\rangle$ and $\overline{C_1} = C_1/\langle f_1\rangle$ to $C_2'/\langle (f_2')^k\rangle = C_2'/\langle f_2'\rangle$.

Now the orientation preserving diffeomorphism $\overline{\varphi} \circ \overline{g}$ of S^3 satisfies $\overline{\varphi} \circ \overline{g}(\overline{K}) = \overline{K}$ and $\overline{\varphi} \circ \overline{g}(\overline{C_1}) = \overline{C_2}$. Thus $\overline{K} \cup \overline{C_1}$ and $\overline{K} \cup \overline{C_2}$ are isotopic. $\square$(Theorem 4.3)

5. Examples

In this section, we present two examples illustrating how the periodic construction works according as the initial knot $\overline{K}$ is fibered or not fibered.

First we apply Corollary 4.1 in the case where $\overline{K}$ is not fibered. In such a case we can choose $\overline{C}$ arbitrarily with $lk(\overline{C}, \overline{K}) \neq 0$ to obtain a knot $K^p_{\overline{C}}$ having properties (1) and (2) in Corollary 4.1.

Let T_n $(n \neq 0, \pm 1)$ be a twist knot illustrated in Figure 5.1.



Figure 5.1. A twist knot T_n

Then T_n is a hyperbolic knot, and since the Alexander polynomial of T_n is not monic, it is not fibered [5, 8.16 Proposition]. Suppose that $n > 1$. Then it follows from [47, 15] that $\pi_1(T_n(r))$ is left-orderable for $r \in (-4n, 4)$. Furthermore, it is known by [45] that $\pi_1(T_n(4))$ is left-orderable. Hence $\pi_1(T_n(r))$ is left-orderable if $r \in (-4n, 4]$.

Example 5.1. Let us take a 2–component link $T_2 \cup \overline{C}$ as in Figure 5.2; $l = lk(\overline{C}, T_2) = 1$. Let p be any integer with $p > 2$. Take the p–fold cyclic branched cover of S^3 branched along $\overline{C}$ to obtain a knot $K^p_{2,\overline{C}}$ which is the preimage of T_2. Then $K^p_{2,\overline{C}}$ is a knot with cyclic period p whose factor knot is T_2, and enjoys the following properties:

1. $K^p_{2,\overline{C}}$ is a hyperbolic knot in S^3.
2. $K^p_{2,\overline{C}}(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$.
3. $K^p_{2,\overline{C}}(r)$ is not an L–space for all $r \in \mathbb{Q}$.
4. $\pi_1(K^p_{2,\overline{C}}(r))$ is left-orderable whenever $r \in (-8p, 4p]$.



Figure 5.2. The twist knot T_2 and an axis $\overline{C}$

Proof. Assertions (1) and (2) follow from Theorem 4.3(2) once we show that $T_2 \cup \overline{C}$ is a hyperbolic link. Since $T_2 \cup \overline{C}$ is a non-split prime alternating link [28, Theorem 1], it is either a torus link or a hyperbolic link [28, Corollary 2]. The former cannot happen, because T_2 is not a torus knot. Hence $T_2 \cup \overline{C}$ is a hyperbolic link as desired. Since T_2 is non-fibered and $\pi_1(T_2(r))$ is left-orderable for $r \in (-8, 4]$, assertions (3) and (4) follow from Corollary 4.1. □(Example 5.1)

Next we apply Corollary 4.1 in the case where $\overline{K}$ is a fibered knot. Let $T_{m,n}$ ($m > n \geq 2$) be a nontrivial torus knot. Then Clay and Watson [9, Theorem 1.4] show that $\pi_1(T_{m,n}(r))$ is left-orderable if $r < mn - m - n$. Hence $\pi_1(T_{3,2}(r))$ is left-orderable if $r < 3 \cdot 2 - 3 - 2 = 1$. For $T_{-3,2}$, since $T_{3,2}(r)$ is orientation reversingly diffeomorphic to $T_{-3,2}(-r)$, we see that $\pi_1(T_{-3,2}(r))$ is left-orderable if $-1 < r$.

Example 5.2. Let us take a 2–component link $T_{-3,2} \cup \overline{C}$ as in Figure 5.3; $l = lk(\overline{C}, T_{-3,2}) = 1$. Let p be any integer with $p > 2$. Take the p–fold cyclic branched cover of S^3 branched along the trivial knot $\overline{C}$ to obtain a knot $K^p_{-3,2,\overline{C}}$ which is the preimage of $T_{-3,2}$. Then $K^p_{-3,2,\overline{C}}$ is a knot with cyclic period p whose factor knot is $T_{-3,2}$.

(1) $K^p_{-3,2,\overline{C}}$ is a hyperbolic knot in S^3.
(2) $K^p_{-3,2,\overline{C}}(r)$ is a hyperbolic 3–manifold for all $r \in \mathbb{Q}$.
(3) $K^p_{-3,2,\overline{C}}(r)$ is not an L–space for all $r \in \mathbb{Q}$.
(4) $\pi_1(K^p_{-3,2,\overline{C}}(r))$ is left-orderable whenever $r \in (-p, \infty)$.



(i) (ii)

FIGURE 5.3. The trefoil knot $T_{-3,2}$ and an unknotted circle $\overline{C}$

Proof of Example 5.2. Assertions (1) and (2) follow from Theorem 4.3(2) once we see that $T_{-3,2} \cup \overline{C}$ is a hyperbolic link. Since as illustrated in Figure 5.3(i) $T_{-3,2} \cup \overline{C}$ is a non-split prime alternating link [28, Theorem 1], it is either a torus link or a hyperbolic link [28, Corollary 2]. If we have the former case, then $T_{-3,2}$ is isotopic to $\overline{C}$ which is a trivial knot, a contradiction. Hence $T_{-3,2} \cup \overline{C}$ is a hyperbolic link as desired.

To see (3) and (4), we apply Corollary 4.1. Since $T_{-3,2}$ is fibered, we need to show that for any fiber surface $\overline{S}$ of $E(T_{-3,2})$, $|\overline{C} \cap \overline{S}|$ is strictly bigger than the algebraic intersection number between $\overline{C}$ and $\overline{S}$, i.e. $lk(\overline{C}, T_{-3,2})$.

In Figure 5.3(ii), we give a minimal genus Seifert surface F of $T_{-3,2}$, which is a once-punctured torus with $\partial F = T_{-3,2}$. Put $\overline{S} = F \cap E(T_{-3,2})$. Then by [12, Lemma 5.1] $\overline{S}$ is a fiber surface of $E(T_{-3,2})$. We see that $|\overline{C} \cap \overline{S}| = 5$ and the algebraic intersection number between $\overline{C}$ and $\overline{S}$ is one. Assume for a contradiction that we have another fiber surface $\overline{S}'$ of $E(T_{-3,2})$ such that $|\overline{C} \cap \overline{S}'| < |\overline{C} \cap \overline{S}|$. Since $\overline{S}$ and $\overline{S}'$ are fiber surfaces of $E(T_{-3,2})$, they are isotopic; see [12, Lemma 5.1], [46]. This then implies that we can isotope $\overline{C}$ to $\overline{C}'$ in $E(T_{-3,2})$ so that $|\overline{C}' \cap \overline{S}| < |\overline{C} \cap \overline{S}|$.

Claim 5.3. *There exists a smooth map φ from a semi-disk D into $E(T_{-3,2})$ such that $\varphi^{-1}(\overline{C})$ is an arc $c \subset \partial D$ and $\varphi^{-1}(\overline{S})$ is the arc $\alpha = \overline{\partial D - c}$.*

Proof of Claim 5.3. Let $\Phi : S^1 \times [0,1] \to E(T_{-3,2})$ be a smooth map giving an isotopy between $\overline{C}(= \Phi(S^1 \times \{0\})$ to $\overline{C}'(= \Phi(S^1 \times \{1\})$. We may assume Φ is transverse to $\overline{S}$. Furthermore, the essentiality of $\overline{S}$ in $E(T_{-3,2})$ enables us to modify Φ to eliminate the circle components as usual. Since $|\overline{C}' \cap \overline{S}| < |\overline{C} \cap \overline{S}| = 5$ and the algebraic intersection number between $\overline{C}'$ and $\overline{S}$ coincides with the algebraic intersection number between $\overline{C}$ and $\overline{S}$, we have $|\overline{C}' \cap \overline{S}| = 1$ or 3. Thus $\Phi^{-1}(\overline{S})$ consists of three properly embedded arcs α, α' and β, where $\partial\alpha \subset S^1 \times \{0\}$, $\partial\alpha' \subset S^1 \times \{0\}$, and β connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$ (Figure 5.4(i), (ii)), consists of four properly embedded arcs α, β, β' and β'', where $\partial\alpha \subset S^1 \times \{0\}$, and each of β, β', β'' connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$ (Figure 5.4(iii)), or consists of four properly embedded arcs α, α', β and γ, where $\partial\alpha \subset S^1 \times \{0\}$, $\partial\alpha' \subset S^1 \times \{0\}$, β connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$, and $\partial\gamma \subset S^1 \times \{1\}$ (Figure 5.4(iv)). In either case there is a semi-disk D cobounded by α and an arc $c \subset S^1 \times \{0\}$.

Putting $\varphi = \Phi|_D : D \to E(T_{-3,2})$, we obtain a required smooth map. □(Claim 5.3)

Cut open $E(T_{-3,2})$ along $\overline{S}$ to obtain a product 3–manifold $\overline{S} \times [0,1]$. The circle $\overline{C}$ is cut into five arcs c_1, c_2, c_3, c_4 and c_5 as in Figure 5.3(ii). Note that $\partial c_1 \subset \overline{S} \times \{0\}$, $\partial c_3 \subset \overline{S} \times \{1\}$, and each of c_2, c_4, c_5 connects $\overline{S} \times \{0\}$ and $\overline{S} \times \{1\}$. Moreover, we see that c_1 and c_3 are linking once relative their boundaries.

On the other hand, since c is either c_1 or c_3, Claim 5.3 shows that c_1 and c_3 are unlinked relative their boundaries. This contradiction shows that for any fiber surface $\overline{S}$, $|\overline{C} \cap \overline{S}| > lk(\overline{C}, T_{-3,2})$ as desired.

Since $\pi_1(T_{-3,2}(r))$ is left-orderable if $r \in (-1, \infty)$, the conclusions (3) and (4) follow from Corollary 4.1. This completes the proof of Example 5.2. □(Example 5.2)



(i) (ii)

(iii) (iv)

Figure 5.4. $\Phi^{-1}(\overline{S})$ in $S^1 \times [0,1]$

For any fibered knot K in S^3 a minimal genus Seifert surface in $E(K)$ is a fiber surface and vice versa, and furthermore, any fiber surface is unique up to isotopy. See [12, Lemma 5.1], [46].

6. Surgeries on alternating knots

Theorem 1.5 in [38], together with [39, Proposition 9.6] ([34, Proof of Corollary 1.3], [18, Claim 2]), shows that for an alternating knot K which is not a $(p,2)$–torus knot, $K(r)$ is not an L–space for all $r \in \mathbb{Q}$.

We say that an alternating knot is *positive* (resp. *negative*) if it has a reduced alternating diagram such that each of the crossings is positive (resp. negative). An alternating knot is *special* if it is either positive or negative.

In [3] Boyer, Gordon and Watson prove:

Proposition 6.1 ([3])**.** *Let K be a prime alternating knot in S^3.*

1. *If K is not a special alternating knot, then $\pi_1(K(\frac{1}{n}))$ is left-orderable for all non-zero integers n.*
2. *If K is a positive* (*resp. negative*) *alternating knot, then $\pi_1(K(\frac{1}{n}))$ is left-orderable for all positive* (*resp. negative*) *integers n.*

For each integer $p > 0$, we apply the periodic construction to produce infinitely many alternating knots K such that $\pi_1(K(\frac{p}{n}))$ is left-orderable for all non-zero integers n.

Let $\overline{K}$ be an alternating knot. For convenience, we position $\overline{K} \subset \mathbb{R}^3 = S^3 - \{\infty\}$ so that $\overline{K}$ lies in the xy–plane except near crossings of $\overline{K}$, where $\overline{K}$ lies on a "bubble" as in [28]. Then we say an unknotted circle $\overline{C} \subset S^3 - \overline{K}$ is *perpendicular* if it passes ∞ and intersects the xy–plane exactly once. Note that $\overline{C} \cap \mathbb{R}^3$ is perpendicular to the xy–plane. See Figure 6.1, in which the dot indicates a perpendicular circle $\overline{C}$.



FIGURE 6.1. An alternating knot $\overline{K}$ and a perpendicular circle $\overline{C}$

Proposition 6.2. *Let $\overline{K}$ be a prime alternating knot in S^3 and $\overline{C}$ a perpendicular circle with $lk(\overline{C}, \overline{K}) = l$. Let p be an integer such that $p \geq 2$ and $(p, l) = 1$. Take the p–fold cyclic branched cover of S^3 branched along $\overline{C}$ to obtain a periodic knot $K^{p}_{\overline{C}}$ which is the preimage of $\overline{K}$. Then we have:*

1. *$K^{p}_{\overline{C}}$ is an alternating knot.*
2. *If $\overline{K}$ is not a special alternating knot, then $\pi_1(K^{p}_{\overline{C}}(\frac{p}{n}))$ is left-orderable for all non-zero integers n.*
3. *If $\overline{K}$ is a positive (resp. negative) alternating knot, then $\pi_1(K^{p}_{\overline{C}}(\frac{p}{n}))$ is left-orderable for all positive (resp. negative) integers n.*

Proof of Proposition 6.2. The first assertion follows immediately from diagramatic consideration. The conclusions (2) and (3) follow from Proposition 6.1 and Theorem 2.1. $\square$(Proposition 6.2)

Remark 6.3. In Proposition 6.2, if $\overline{K}$ is not a $(p, 2)$–torus knot, then $K^{p}_{\overline{C}}$ is not a $(p', 2)$–torus knot. For otherwise, the argument in the proof of Lemma 2.3 implies that $\overline{K}$ is a torus knot. Since it is alternating, it is a $(p, 2)$–torus knot for some odd integer q [32, Theorem 3.2], a contradiction. Therefore, as mentioned in the beginning of this section, $K^{p}_{\overline{C}}(r)$ is not an L–space for all $r \in \mathbb{Q}$.

Applying Proposition 6.2 and Remark 6.3, we have:

Example 6.4. Take an alternating knot $\overline{K}$ and a perpendicular circle $\overline{C}$ as illustrated in Figure 6.1; $lk(\overline{C}, \overline{K}) = 1$. Note that $\overline{K}$ is not a special alternating knot.

Hence for any integer $p \geq 2$, $K_{\overline{C}}^{p}$ is an alternating knot, $K_{\overline{C}}^{p}(r)$ is not an L–spaces for all $r \in \mathbb{Q}$, and $\pi_1(K_{\overline{C}}^{p}(\frac{p}{n}))$ is left-orderable for all non-zero integers n.

7. Proofs of Theorems 1.2 and 1.3.

The goal of this section is to prove Theorems 1.2 and 1.3.

Proof of Theorem 1.2. Let us consider the connected sum $T_{-3,2} \sharp T_{3,2}$. We recall the following well-known general fact.

Claim 7.1. *Let $K_1, \dots, K_n$ be nontrivial knots. Then $(K_1 \sharp \cdots \sharp K_n)(r)$ is irreducible for all $r \in \mathbb{Q}$.*

Proof of Claim 7.1. First we note that the exterior $E(K_1 \sharp \cdots \sharp K_n)$ is a union of a composing space C_n (i.e. [disk with $n-$holes]$\times S^1$) and $E(K_1), \dots, E(K_n)$. Hence for any $r \in \mathbb{Q}$, $(K_1 \sharp \cdots \sharp K_n)(r)$ is a union of $C_n \cup V$ and $E(K_1), \dots, E(K_n)$, where V is a filled solid torus. Note that $C_n \cup V$ has a Seifert fibration over the disk with $(n-1)$–holes with at most one exceptional fiber, and hence it is irreducible and boundary-irreducible. Then since $C_n \cup V$ and $E(K_i)$ $(1 \leq i \leq n)$ are irreducible and boundary-irreducible, $(K_1 \sharp \cdots \sharp K_n)(r)$ is also irreducible. □(Claim 7.1)

Let us regard $T_{-3,2} \sharp T_{3,2}$ as a satellite knot with the companion knot $T_{3,2}$ and the pattern knot $T_{-3,2}$. Since $\pi_1(T_{-3,2}(r))$ is left-orderable if $r > -1$ [9], and $(T_{-3,2} \sharp T_{3,2})(r)$ is irreducible for all $r \in \mathbb{Q}$ (Claim 7.1), Proposition 4.1 in [9] shows that $\pi_1((T_{-3,2} \sharp T_{3,2})(r))$ is also left-orderable if $r > -1$. Using the amphicheirality of $T_{-3,2} \sharp T_{3,2}$, we see that $\pi_1((T_{-3,2} \sharp T_{3,2})(r))$ is left-orderable also when $r < 1$. Therefore it is left-orderable for all $r \in \mathbb{Q}$. Note that $T_{-3,2} \sharp T_{3,2}$ is a fibered knot.

Before we apply Corollary 4.1, for ease of handling, take the connected sum $(T_{-3,2} \sharp T_{3,2}) \sharp T_2$. The Alexander polynomial of $(T_{-3,2} \sharp T_{3,2}) \sharp T_2$ is $(t^2-t+1)^2(2t^2-5t+2)$, which is not monic, and hence $(T_{-3,2} \sharp T_{3,2}) \sharp T_2$ is not fibered. We regard $(T_{-3,2} \sharp T_{3,2}) \sharp T_2$ as a satellite knot with the companion knot T_2 and the pattern knot $T_{-3,2} \sharp T_{3,2}$. As we observe above, $\pi_1((T_{-3,2} \sharp T_{3,2})(r))$ is left-orderable for all $r \in \mathbb{Q}$. Moreover by Claim 7.1 $(T_{-3,2} \sharp T_{3,2} \sharp T_2)(r)$ is irreducible for all $r \in \mathbb{Q}$. We apply [9, Proposition 4.1] again to conclude that $(T_{-3,2} \sharp T_{3,2} \sharp T_2)(r)$ has the left-orderable fundamental group for all $r \in \mathbb{Q}$.

To obtain hyperbolic knots with this property, we will apply the periodic construction (Corollary 4.1). Let us put $\overline{K} = T_{-3,2} \sharp T_{3,2} \sharp T_2$ and take an unknotted circle $\overline{C}$ as in Figure 7.1; $lk(\overline{C}, \overline{K}) = 1$.

Since $\overline{K} \cup \overline{C}$ is a non-split prime alternating link [28, Theorem 1], it is either a torus link or a hyperbolic link [28, Corollary 2]. The former cannot happen, because $\overline{K}$ is not a torus knot. Hence $\overline{K} \cup \overline{C}$ is a hyperbolic link. Let $p > 2$ be any integer.


$\overline{\mathrm{C}}$
$\overline{\mathrm{K}} = \mathrm{T}_{-3,2}$ # $\mathrm{T}_{3,2}$ # T_2

Figure 7.1. $\overline{K} \cup \overline{C}$

Take the p–fold cyclic branched cover of S^3 branched along $\overline{C}$ to obtain a periodic knot $K_{\overline{C}}^{p}$ which is the preimage of $\overline{K}$.

It follows from Corollary 4.1 and Theorem 4.3(2) that $K_{\overline{C}}^{p}$ is a hyperbolic knot and enjoys the properties (1), (2) and (3) in Theorem 1.2. By changing p, we obtain infinitely many such knots. For instance, see Remark 4.2. □(Theorem 1.2)

Remark 7.2.

(1) By Theorem 4.3 there are infinitely many unknotted circles for $\overline{K} = T_{-3,2} \sharp T_{3,2} \sharp T_2$, and for each unknotted circle $\overline{C}$ we obtain infinitely many hyperbolic knots $K_{\overline{C}}^{p}$, where p and $lk(\overline{C}, \overline{K})$ are relatively prime.

(2) Recall that any knot K obtained by the "periodic construction", for instance a knot obtained in the proof of Theorem 1.2, is not fibered and every nontrivial surgery on K is a left-orderable, non–L–space surgery. So we can apply Corollary 4.1 again to the knot K and an arbitrarily chosen unknotted circle to obtain yet further infinitely many non-fibered knots K' each of which has the (same) factor knot K. Then r–surgery on K' is also a left-orderable, non–L–space surgery for all $r \in \mathbb{Q}$. We can apply this procedure repeatedly arbitrarily many times.

(3) Let K be the knot 10_{99} in Rolfsen's knot table [42]. Recently Adam Clay [6] uses an epimorphism from $E(K)$ to $E(T_{3,2})$ which preserves the peripheral subgroup [23] to show that every nontrivial surgery on K is left-orderable surgery. Since K has no cyclic period [22, Appendix F], this curious example cannot be explained by the periodic construction.

The *tunnel number* of a knot K, denoted by $t(K)$, is defined as the least number of mutually disjoint arcs t_i $(i = 1, \dots, n)$ with $t_i \cap K = \partial t_i$ such that the exterior of the regular neighborhood of $K \cup t_1 \cup \cdots \cup t_n$ is a handlebody. Given a 3–manifold M, we denote its Heegaard genus by $g(M)$.

Proof of Theorem 1.3. Let us take the knot $K_{\overline{C}}^{4}$ given in the proof of Theorem 1.2. As mentioned in the proof of Theorem 1.2, $T_{-3,2} \sharp T_{3,2} \sharp T_2$ is not fibered, hence

$K_{\frac{4}{C}}$ is not fibered; see the proof of Proposition 3.1. Furthermore, it follows from Theorem 1.2 that $\pi_1(K_{\frac{4}{C}}(r))$ is left-orderable for all $r \in \mathbb{Q}$.

Take an unknotted circle $\overline{C}_\mu$ which is a meridian of $K_{\frac{4}{C}}$, and apply Corollary 4.1 to the pair $K_{\frac{4}{C}}$ and $\overline{C}_\mu$. Then we obtain the connected sum of p–copies of $K_{\frac{4}{C}}$, denoted by K_p, which satisfies the conclusions (3) and (4) in Theorem 1.3.

It remains to prove (1) and (2) in Theorem 1.3. Since $g(T_{-3,2} \sharp T_{3,2} \sharp T_2) = 3$, by Remark 4.2 $g(K_{\frac{4}{C}}) \geq 4 \cdot 3 = 12$. Thus $g(K_p) = pg(K_{\frac{4}{C}}) \geq 12p$. Concerning the tunnel number $t(K_p)$, it follows from [43] that $t(K_p) \geq p$. This then implies $g(E(K_p)) = t(K_p)+1 \geq p+1$. Now [41, Corollary 4.4 and Remark in p.627] asserts that $K_p(r)$ has Heegaard genus at least p for all but finitely many $r \in \mathbb{Q}$ once the claim below holds.

Claim 7.3. *There is no Möbius band properly embedded in $E(K_p)$.*

Proof of Claim 7.3. The proof of Claim 7.1 shows that $K_p(r)$ is a Haken 3–manifold for all $r \in \mathbb{Q}$. Now suppose for a contradiction that $E(K_p)$ contains a properly embedded Möbius band F. Then performing Dehn surgery along the boundary slope r' of F, we obtain a 3–manifold $K_p(r')$ containing a projective plane. This then implies that $K_p(r')$ is $\mathbb{R}P^3$ or a reducible manifold having $\mathbb{R}P^3$ as a connected summand. In either case, we have a contradiction. Therefore there is no Möbius band properly embedded in $E(K_p)$ as claimed. □(Claim 7.3)

For the given integer N, taking $p > N$, the assertions (1) and (2) hold for $K = K_p$. This completes a proof of Theorem 1.3. □(Theorem 1.3)

8. QUESTIONS

Let K be a knot in S^3. Define the set of left-orderable slopes for K as

$$\mathcal{S}_{LO}(K) = \{r \in \mathbb{Q} \mid \pi_1(K(r)) \text{ is left-orderable}\}.$$

For any knot K, $H_1(K(0)) \cong \mathbb{Z}$, and hence $\{0\} \subset \mathcal{S}_{LO}(K)$. If K is the trivial knot then $\mathcal{S}_{LO}(K) = \{0\}$. In the present note, we demonstrate that there are infinitely many hyperbolic knots K with $\mathcal{S}_{LO}(K) = \mathbb{Q}$.

It seems interesting to ask:

Question 8.1. If K is a nontrivial knot in S^3, then does $\mathcal{S}_{LO}(K)$ contain $(-1,1)\cap\mathbb{Q}$?

Recently Li and Roberts [24, Corollary 1.2] prove that for any hyperbolic knot K, there exists a constant N_K such that $\{\frac{1}{n} \mid |n| > N_K\} \subset \mathcal{S}_{LO}(K)$.

More strongly we would like to ask:

Question 8.2. If K is a nontrivial knot in S^3, then does $\mathcal{S}_{LO}(K)$ contain $(-\infty,1)\cap\mathbb{Q}$ or $(1,\infty)\cap\mathbb{Q}$?

The argument in the proof of [9, Theorem 1.4] shows that $\mathcal{S}_{LO}(T_{3,2}) = (-\infty, 1) \cap \mathbb{Q}$ and $\mathcal{S}_{LO}(T_{-3,2}) = (-1, \infty) \cap \mathbb{Q}$.

Question 8.3. If $\mathcal{S}_{LO}(K) = (-\infty, 1) \cap \mathbb{Q}$ or $\mathcal{S}_{LO}(K) = (-1, \infty) \cap \mathbb{Q}$, then is K a trefoil knot $T_{3,2}$ or $T_{-3,2}$, respectively?

Acknowledgements – We would like to thank Adam Clay for informing us his curious example mentioned in Remark 7.2(3).

References

[1] I. Aitchison; Isotoping and twisting knots and ribbons, Thesis–University of California, Berkeley May 1984.

[2] M. Boileau and B. Zimmermann; The π–orbifold group of a link, Math. Z. **200** (1989), 187–208.

[3] S. Boyer, C. McA. Gordon and L. Watson; On L–spaces and left-orderable fundamental groups, to appear in Math. Ann..

[4] S. Boyer, D. Rolfsen and B. Wiest; Orderable 3-manifold groups, Ann. Inst. Fourier **55** (2005), 243–288.

[5] G. Burde and H. Zieschang; Knots, de Gruyter Studies in Mathematics, 5. Walter de Gruyter & Co., Berlin, 2003.

[6] A. Clay; private communication.

[7] A. Clay, T. Lidman and L. Watson; Graph manifolds, left-orderability and amalgamation, preprint.

[8] A. Clay and M. Teragaito; Left-orderability and exceptional Dehn surgery on two-bridge knots, to appear in Contemp. Math. Amer. Math. Soc.

[9] A. Clay and L. Watson; On cabled knots, Dehn surgery, and left-orderable fundamental groups, Math. Res. Lett. **18** (2011), 1085–1095.

[10] A. Clay and L. Watson; Left-orderable fundamental groups and Dehn surgery, International Mathematics Research Notices, published online: 2012.

[11] A. Edmonds; Least area Seifert surfaces and periodic knots, Topology Appl. **18** (1984) 109–113.

[12] A. Edmonds and C. Livingston; Group actions on fibered three-manifolds, Comment. Math. Helv. **58** (1983) 529–542.

[13] J. Greene; Alternating links and left-orderability, preprint.

[14] R. Hakamata and M. Teragaito; Left-orderable fundamental group and Dehn surgery on twist knots, preprint.

[15] R. Hakamata and M. Teragaito; Left-orderable fundamental group and Dehn surgery on genus one two-bridge knots, preprint.

[16] C. Hayashi and K. Motegi; Dehn surgery on knots in solid tori creating essential annuli, Trans. Amer. Math. Soc. **349** (1997), 4897–4930.

[17] C. Hayashi and K. Shimokawa; Symmetric knots satisfy the cabling conjecture, Math. Proc. Camb. Phil. Soc. **123** (1998) 501-529.

[18] K. Ichihara and I.D. Jong; Cyclic and finite surgeries on Montesinos knots, Algebr. Geom. Topol. **9** (2009), 731?742.

[19] T. Ito; Non-left-orderable double branched coverings, preprint.

[20] W. Jaco and P. Shalen; Seifert fibered spaces in 3–manifolds, Mem. Amer. Math. Soc. **21** (1979), no. 220, viii+192.

[21] K. Johannson; Homotopy equivalences of 3–manifolds with boundaries, Lect. Notes in Math. vol. 761, Springer-Verlag, 1979.

[22] A. Kawauchi; A survey of knot theory, BirkhNauser-Verlag, Basel, Boston, and Berlin, 1996

[23] T. Kitano and M. Suzuki, A partial order in the knot table, Exp. Math. **14** (2005), 385–390.

[24] T. Li and R. Roberts; Taut foliations in knot complements, preprint.

[25] E. Luft and X. Zhang; Cabling conjecture for symmetric knots, Math. Ann. **298** (1994) 489–496.

[26] W. Meeks and P. Scott; Finite group actions on 3-manifolds, Invent. Math. **86** (1986) 287–346.
[27] W. Meeks and S.-T. Yau; The equivariant Dehn's lemma and loop theorem, Comment. Math. Helv. **56** (1981) 225–239.
[28] W. Menasco; Closed incompressible surfaces in alternating knots and link complements, Topology **23** (1984) 37–44.
[29] K. Miyazaki and K. Motegi; Seifert fibered manifolds and Dehn surgery III, Comm. Anal. Geom. 7 (1999), 551–582.
[30] K. Miyazaki and K. Motegi; Toroidal surgery on periodic knots, Pacific J. Math. **193** (2000) 381–396.
[31] J. Morgan and H. Bass (eds.); The Smith conjecture, Academic Press, 1984.
[32] K. Murasugi; On the Alexander polynomial of the alternating knot, Osaka Math. J. **10** (1958) 181–189.
[33] S. Naik; Periodicity, genera and Alexander polynomials of knots, Pacific J. Math. **166** (1994) 357–371.
[34] Y. Ni; Knot Floer homology detects fibred knots, Invent. Math. **170** (2007) 577–608.
[35] Y. Ni; Erratum: Knot Floer homology detects fibred knots, Invent. Math. **177** (2009), 235–238.
[36] P. Ozsváth and Z. Szabó; Holomorphic disks and topological invariants for closed three-manifolds, Ann. of Math. **159** (2004), 1027–1158.
[37] P. Ozsváth and Z. Szabó; Holomorphic disks and three-manifold invariants: properties and applications, Ann. of Math. **159** (2004), 1159–1245.
[38] P. Ozsváth and Z. Szabó; On knot Floer homology and lens space surgeries, Topology **44** (2005), 1281–1300.
[39] P. Ozsváth and Z. Szabó; Knot Floer homology and rational surgeries, Algebr. Geom. Topol. **11** (2011), 1–68.
[40] T. Peters; On L-spaces and non left-orderable 3-manifold groups, preprint.
[41] Y. Rieck; Heegaard structures of manifolds in the Dehn filling spaces, Topology **39** (2000) 619–641.
[42] D. Rolfsen; Knots and links, Publish or Perish, 1976.
[43] M. Scharlemann and J. Schultens; The tunnel number of the connect sum of n knots is at least n, Topology **38** (1999) 265–270.
[44] P. A. Smith; Transformations of finite period. II, Ann. of Math. **40** (1939) 690?711.
[45] M. Teragaito; Left-orderability and exceptional Dehn surgery on twist knots, to appear in Canad. Math. Bull.
[46] W. Thurston; A norm for the homology of 3-manifolds, Mem. Amer. Math. Soc. **59** (1986), no. 339, i–vi, 99–130.
[47] A. Tran; On left-orderable fundamental groups and Dehn surgeries on twist knots, preprint.
[48] A. Tran; On left-orderable fundamental groups and Dehn surgeries on double twist knots, preprint.

Department of Mathematics, Nihon University, Tokyo 156–8550, Japan
E-mail address: `motegi@math.chs.nihon-u.ac.jp`

Department of Mathematics and Mathematics Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-Hiroshima, 739–8524, Japan
E-mail address: `teragai@hiroshima-u.ac.jp`